

University of Edinburgh

**A NOTE ON ULTRAVIOLET SUPPRESSED QUASI-OPTIMAL
DOMAIN WALL FERMIONS**

ARTAN BORIÇI

ABSTRACT. In a recent work Chiu proposed to modify domain wall fermions that allow in an optimal way fewer number of flavors than in the standard case. This is done using a variant of domain wall fermions, the so-called truncated overlap fermions. In this note I discuss the possibility to implement his proposal for the original variant of domain wall fermions. I make also some remarks on dynamical simulations with ultraviolet suppressed domain wall fermions.

Domain wall and overlap fermions. I will briefly review the definitions of domain wall fermions [Kaplan, 1992, Furman and Shamir, 1995]. Let N be the size of the extra dimension, D_W the Wilson-Dirac operator, and m the bare fermion mass. Then, the domain wall fermion action is given by:

$$(0.1) \quad S_{DWF} := \bar{\Psi} \mathcal{D} \Psi = \sum_{i=1}^N \bar{\psi}_i [a_5(D^{\parallel} - \mathbb{1})\psi_i + P_+ \psi_{i+1} + P_- \psi_{i-1}]$$

together with boundary conditions:

$$(0.2) \quad \begin{aligned} P_+(\psi_{N+1} + m\psi_1) &= 0 \\ P_-(\psi_0 + m\psi_N) &= 0 \end{aligned}$$

Here $\mathcal{M}$ is the five-dimensional fermion matrix of the theory and $D^{\parallel} = M - D_W$ with $M \in (0, 2)$ being a mass parameter. The alternative theory of chiral fermions on the lattice is formulated by [Narayanan and Neuberger, 1993]. In the domain wall setting this theory can be equivalently formulated using truncated overlap fermions [Boriçi, 2000a, Boriçi, 2000b]:

$$(0.3) \quad S_{TOV} := \bar{\Psi} \mathcal{D}_{TOV} \Psi = \sum_{i=1}^N \bar{\psi}_i [a_5(D^{\parallel} - \mathbb{1})\psi_i + (a_5 D^{\parallel} + \mathbb{1})P_+ \psi_{i+1} + (a_5 D^{\parallel} + \mathbb{1})P_- \psi_{i-1}]$$

with the same boundary conditions as above. In both cases the four dimensional lattice spacing is set to one.

It is well-known that domain wall fermions in both variants can be dimensionally reduced to give an effective theory of four dimensional chiral fermions [Neuberger, 1998b, Kikukawa and Noguchi, 1999, Boriçi, 1999a]. The Dirac operator of the theory is given by the Neuberger overlap operator [Neuberger, 1998a]:

$$(0.4) \quad D = \frac{1+m}{2} \mathbb{1} - \frac{1-m}{2} \gamma_5 \operatorname{sgn}(H)$$

where $H = \gamma_5 D^{\parallel}$ in case of truncated overlap fermions with transfer matrix [Boriçi, 2000a]:

$$(0.5) \quad T = \frac{\mathbb{1} + a_5 H}{\mathbb{1} - a_5 H}$$

For domain wall fermions it was shown that [Boriçi, 1999a]:

$$(0.6) \quad \mathcal{H} = \gamma_5 \frac{D^{\parallel}}{2 - a_5 D^{\parallel}} = H \frac{1}{2 - a_5 D^{\parallel}}$$

Recently, Chiu proposed to use a transfer matrix which depends on the 5D time slice [Chiu, 2002]:

$$(0.7) \quad T_i = \frac{\mathbb{1} + \omega_i a_5 H}{\mathbb{1} - \omega_i a_5 H}$$

where $\omega_i, i = 1, \dots, N$ are positive coefficients and $a_5 = 1$. These coefficients are chosen such as to correspond to the optimal Chebyshev rational approximation of the sign function, the Zolotarev approximation [Zolotarev, 1877]. In his paper, Chiu showed how to implement this idea in case of truncated overlap fermions. Below I discuss Chiu's idea in the case of domain wall fermions.

Quasi-optimal domain wall fermions. I postulate the optimal domain wall fermion action given by:

$$(0.8) \quad \bar{\Psi} \mathcal{D} \Psi = \sum_{i=1}^N \bar{\psi}_i [(\omega_i a_5 D^{\parallel} - \mathbb{1}) \psi_i + P_+ \psi_{i+1} + P_- \psi_{i-1}]$$

To show that such domain wall fermions are optimal it is sufficient to calculate tranfer matrices $T_i, i = 1, \dots, N$ from $\mathcal{M}$ and find that they are in the optimal form (0.7). The fermion matrix can be written as an $N \times N$ block partition along the fifth dimension:

$$\mathcal{M} = \begin{pmatrix} \omega_1 a_5 D^{\parallel} - \mathbb{1} & P_+ & & -mP_- \\ P_- & \omega_2 a_5 D^{\parallel} - \mathbb{1} & \ddots & \\ & \ddots & \ddots & P_+ \\ -mP_+ & & P_- & \omega_N a_5 D^{\parallel} - \mathbb{1} \end{pmatrix}$$

Multiplying from the right by the permutation matrix:

$$\begin{pmatrix} P_+ & P_- & & \\ & P_+ & \ddots & \\ & & \ddots & P_- \\ P_- & & & P_+ \end{pmatrix}$$

I obtain:

$$\gamma_5 \begin{pmatrix} (\omega_1 a_5 H P_+ - \mathbb{1})(P_+ - mP_-) & \omega_1 a_5 H P_- + \mathbb{1} & & \\ & \omega_2 a_5 H P_+ - \mathbb{1} & \ddots & \\ & & \ddots & \omega_{N-1} a_5 H P_- + \mathbb{1} \\ (\omega_N a_5 H P_- + \mathbb{1})(P_- - mP_+) & & & \omega_N a_5 H P_+ - \mathbb{1} \end{pmatrix}$$

Further, multiplying this result from the left by the inverse of the diagonal matrix:

$$\begin{pmatrix} \omega_1 a_5 H P_+ - \mathbb{1} & & & \\ & \omega_2 a_5 H P_+ - \mathbb{1} & & \\ & & \ddots & \\ & & & \omega_N a_5 H P_+ - \mathbb{1} \end{pmatrix}$$

I get:

$$(0.9) \quad \mathcal{T}(m) := \begin{pmatrix} P_+ - mP_- & -T_1 & & \\ & \mathbb{1} & \ddots & \\ & & \ddots & -T_{N-1} \\ -T_N(P_- - mP_+) & & & \mathbb{1} \end{pmatrix}$$

with transfer matrices being defined by:

$$T_i = \frac{\mathbb{1}}{\mathbb{1} - \omega_i a_5 H P_+} (\mathbb{1} + \omega_i a_5 H P_-), \quad i = 1, \dots, N$$

By requiring transfer matrices being in the form:

$$T_i = \frac{\mathbb{1} + \omega_i a_5 \mathcal{H}}{\mathbb{1} - \omega_i a_5 \mathcal{H}}$$

it is easy to see that [Boriçi, 1999a]:

$$(0.10) \quad \mathcal{H} = H \frac{1}{2 - \omega_i a_5 D \parallel}$$

Note that $\mathcal{H}$ depends in the time slice i . By ignoring the denominator the resulting transfer matrix will be of the optimal form (0.7). This may lead to suboptimal approximation of the sign function and must be studied numerically.

Finally to derive the four dimensional Dirac operator one has to compute:

$$\det D^{(N)} = \frac{\det \mathcal{T}(m)}{\det \mathcal{T}(1)}$$

It is easy to show that the determinant of the $N \times N$ block matrix $\mathcal{T}(m)$ is given by:

$$\det \mathcal{T}(m) = \det[(P_+ - mP_-) - T_1 \cdots T_N(P_- - mP_+)]$$

or

$$\det \mathcal{T}(m) = \det\left[\frac{1+m}{2}\gamma_5(\mathbb{1} + T_1 \cdots T_N) + \frac{1-m}{2}(\mathbb{1} - T_1 \cdots T_N)\right]$$

which gives:

$$D^{(N)} = \frac{1+m}{2}\mathbb{1} + \frac{1-m}{2}\gamma_5 \frac{\mathbb{1} - T_1 \cdots T_N}{\mathbb{1} + T_1 \cdots T_N}$$

In the large N limit one gets the Neuberger operator (0.4).

Ultraviolet suppressed domain wall fermions. One difficulty of dynamical simulations with domain wall fermions is the presence of many heavy fermion modes which are simulated together with the interesting light fermion. The problem, generally known as the ultraviolet slowing down becomes more severe in this case. A possible solution is to subtract the contribution of heavy modes in the first place [Izubuchi, RBC Collaboration, 2002].

Here I discuss another solution which is based on the recent proposal of ultraviolet suppressed fermions [Boriçi, 2002a]. This proposal simply states that the lattice Dirac operator can be modified to give:

$$(0.11) \quad D = \frac{\mu}{a_5} \Gamma_5 \tanh \frac{a_5 \Gamma_5 D_{\text{DWF}}}{\mu}$$

where D_{DWF} is the input domain wall operator, a the lattice spacing and $\mu > 0$ is a dimensionless parameter and $\Gamma_5 = \gamma_5 R_5$. R_5 is the reflection operator with respect to the 5-dimensional midpoint [Blum *et al*, 2000]. The Dirac operator converges to the input operator in the continuum limit and is local and unitary as shown in detail in [Boriçi, 2002a]. The input theory is also recovered in the limit $\mu \rightarrow \infty$. For $\mu \rightarrow 0$ one has $D \rightarrow \mu$, i.e. a quenched theory.

Note that the parameter μ has a special role in this theory. Fixing μ at small values, typically $\mu \sim 1$ one can achieve the desired suppression of heavy fermion modes of the input domain wall operator. This way, one localizes the interesting light mode.

At the first sight, it looks that the introduction of a matrix function can lead to potentially difficult computations. In fact, it can be shown that using Lanczos based methods and multi-shift conjugate gradient iterations the computational overhead is small [1].

The benefit using the ultraviolet suppressed operator is hence twofold: one stays within the conventional domain wall formalism of chiral fermions and accelerates dynamical simulations by suppressing both heavy and ultraviolet modes.

End note. This work has been prepared during October 2002 when additions in the paper of Chiu2002 had not already appeared. It is nice to observe that his numerical computations support the proposal of using quasi-optimal domain wall fermions.

REFERENCES

- [Blum *et al*, 2000] T. Blum *et al*, *Quenched Lattice QCD with Domain Wall Fermions and the Chiral Limit*, **hep-lat/0007038**.
- [Boriçi, 1999a] A. Boriçi, *Truncated Overlap Fermions: the link between Overlap and Domain Wall Fermions*, in V. Mitrjushkin and G. Schierholz (eds.), *Lattice Fermions and Structure of the Vacuum*, Kluwer Academic Publishers, 2000.
- [Boriçi, 2000a] A. Boriçi, *Truncated Overlap Fermions*, Nucl. Phys. Proc. Suppl. 83 (2000) 771-773
- [Boriçi, 2000b] A. Boriçi, *Chiral Fermions and Multigrid*, Phys. Rev. D62 (2000) 017505
- [Boriçi, 2002a] A. Boriçi, *Lattice QCD with Suppressed High Momentum Modes of the Dirac Operator*, **hep-lat/0205011**.
- [1] A. Boriçi, *Computational Methods for UV-Suppressed Fermions*, **hep-lat/0208034**.
- [Chiu, 2002] T.W. Chiu, *Optimal domain-wall fermions*, **hep-lat/0209153**
- [Furman and Shamir, 1995] V. Furman, Y. Shamir, *Axial symmetries in lattice QCD with Kaplan fermions*, Nucl. Phys. B439 (1995) 54-78
- [Izubuchi, RBC Collaboration, 2002] T. Izubuchi, RBC Collaboration, *Lattice QCD with dynamical domain wall quarks*, **hep-lat/0210011**
- [Kaplan, 1992] D.B. Kaplan, Phys. Lett. B 228 (1992) 342.
- [Kikukawa and Noguchi, 1999] Y. Kikukawa and T. Noguchi, *Low energy effective action of domain-wall fermion and the Ginsparg-Wilson relation*, Nucl.Phys.Proc.Suppl. 83 (2000) 630-632
- [Narayanan and Neuberger, 1993] R. Narayanan, H. Neuberger, Phys. Lett. B 302 (1993) 62, Nucl. Phys. B 443 (1995) 305.
- [Neuberger, 1998a] H. Neuberger, Phys. Lett. B 417 (1998) 141,
- [Neuberger, 1998b] H. Neuberger, Phys. Rev. D 57 (1998) 5417,
- [Zolotarev, 1877] E. I. Zolotarev, *Application of the elliptic functions to the problems on the functions of least and most deviation from zero*, Zapiskah Rossijskoi Akad. Nauk. (1877). In Russian.